


**LIMIT LAWS**

- $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$
- $\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0$
- $\lim_{x \rightarrow 0} \frac{\sin^2 x}{x} = 0$
- $\lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0$
- $\lim_{x \rightarrow 0} \frac{\sin(ax)}{(bx)} = \lim_{x \rightarrow 0} \frac{\sin(ax)}{\sin(ax)} = \lim_{x \rightarrow 0} \frac{(bx)}{\sin(ax)} = \frac{a}{b}$
- $\lim_{x \rightarrow a} f(x) = L$  (exists) If and only if  $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$
- $f(x)$  is cont at  $a$  if  $\lim_{x \rightarrow a} f(x) = f(a)$

8. **Continuity at a** if  $\lim_{x \rightarrow a^-} f(x) = f(a) = \lim_{x \rightarrow a^+} f(x)$

**TYPES OF DISCONTINUITY**



**ARITHMETIC OF INFINITY**

$1. \infty + \infty = \infty$ $2. n + \infty = \infty$ $3. \infty + 0 = \infty$ $4. 0 + \infty = \infty$	(+)	$1. \infty \cdot \infty = \infty$ $2. n \cdot \infty = \infty$ $3. 0 \cdot \infty = 0$	(x)
$1. \infty - \infty = und$ $2. n - \infty = -\infty$ $3. \infty - n = \infty$ $4. n - n^- = 0^- = -\frac{1}{\infty}$ $5. n^+ - n = 0^+ = \frac{1}{\infty}$	(-)	$1. \infty / \infty = und$ $2. n / \pm \infty = 0$ $3. \infty / n = \infty$ $4. n / 0 = \pm \infty$ $5. n / 0^+ = \infty$ $6. n / 0^- = -\infty$	(÷)
$1. \infty^\infty = \infty$ $2. \infty^n = \infty$ $3. \infty^0 = 1$ $5. 0^\infty = 0$ $4. n^\infty = \infty$	power	$1. -1 \leq \sin(\pm \infty) \leq 1$ $2. -1 \leq \cos(\pm \infty) \leq 1$	Trig

LIMITS(1)

**Basic Rules**

- $\frac{d}{dx} c = 0$  (Constant)
- $\frac{d}{dx} c[f(x)] = c \frac{d}{dx} f(x)$  (constant multiple)
- $\frac{d}{dx} x^n = nx^{n-1}$  (power)
- $\frac{d}{dx} (u \pm v) = \frac{d}{dx} u \pm \frac{d}{dx} v$  (Sum & Difference)
- $\frac{d}{dx} (uv) = v \frac{d}{dx} u + u \frac{d}{dx} v$  (Product)
- $\frac{d}{dx} \left(\frac{u}{v}\right) = \frac{v \frac{d}{dx} u - u \frac{d}{dx} v}{v^2}$  (Quotient)
- $(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$  (Inverse)
- $\frac{d}{dx} f(g(x)) = \frac{d}{dx} f(u) \cdot \frac{d}{dx} g(x)$  where  $u = g(x)$  (Chain Rule)

**Trigonometric Functions**

- $\frac{d}{dx} \sin u = \cos u \frac{du}{dx}$
- $\frac{d}{dx} \cos u = -\sin u \frac{du}{dx}$
- $\frac{d}{dx} \tan u = \sec^2 u \frac{du}{dx}$
- $\frac{d}{dx} \cot u = -\csc^2 u \frac{du}{dx}$
- $\frac{d}{dx} \sec u = \sec u \tan u \frac{du}{dx}$
- $\frac{d}{dx} \csc u = -\csc u \cot u \frac{du}{dx}$

**Inverse Trigonometric Functions**

- $\frac{d}{dx} \sin^{-1} u = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}$
- $\frac{d}{dx} \cos^{-1} u = \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx}$
- $\frac{d}{dx} \tan^{-1} u = \frac{1}{1+u^2} \frac{du}{dx}$
- $\frac{d}{dx} \cot^{-1} u = \frac{-1}{1+u^2} \frac{du}{dx}$
- $\frac{d}{dx} \sec^{-1} u = \frac{1}{|u|\sqrt{u^2-1}} \frac{du}{dx}$
- $\frac{d}{dx} \csc^{-1} u = \frac{-1}{|u|\sqrt{u^2-1}} \frac{du}{dx}$


DERIVATIVES(2)

**Exponential and Logarithmic Functions**

- $\frac{d}{dx} \ln u = \frac{1}{u} \frac{du}{dx}$
- $\frac{d}{dx} \log_a u = \frac{1}{u \ln a} \frac{du}{dx}$
- $\frac{d}{dx} e^u = e^u \frac{du}{dx}$
- $\frac{d}{dx} a^u = a^u \ln a \frac{du}{dx}$
- $\frac{d}{dx} \sqrt{u} = \frac{1}{2\sqrt{u}} \frac{du}{dx}$
- $\frac{d}{dx} |u| = \frac{u}{|u|} \frac{du}{dx}$

WWW.MATHGOTSERVED.COM

**WHERE THE DERIVATIVE DOES NOT EXIST**



Linearization:  $L(x) = f(a) + f'(a)(x-a)$

30. **Mean Value Theorem:** If  $f$  is cont on  $[a,b]$  on and diff on  $(a,b)$

$\Rightarrow$  exist a  $c \in (a,b)$  s.t.  $f'(c) = \frac{f(b)-f(a)}{b-a}$

31. **Rolle's Theorem:** MVT where  $f'(c) = \frac{f(b)-f(a)}{b-a} = 0$

32. **Intermediate Value Theorem:** If  $f$  is cont on  $[a,b]$  and  $d \in [f(a), f(b)]$  then there is a  $c \in [a,b]$  st  $f(c) = d$

**Definition of Derivative**

33.  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

34.  $f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$

35.  $f'(x) \approx$  Average rate of change  
 = Slope of Secant line  
 $= \frac{f(b) - f(a)}{b - a}$

**APPROXIMATING AREA**

- LRAM**<sub>n</sub> =  $w(f(x_1) + f(x_2) + \dots + f(x_{n-1}))$  or  $w_1 f(x_1) + w_2 f(x_2) + \dots + w_{n-1} f(x_{n-1})$
- RRAM**<sub>n</sub> =  $w(f(x_2) + f(x_3) + \dots + f(x_n))$  or  $w_1 f(x_2) + w_2 f(x_3) + \dots + w_{n-1} f(x_n)$
- MRAM**<sub>n</sub> =  $w \left( f\left(\frac{x_1+x_2}{2}\right) + f\left(\frac{x_2+x_3}{2}\right) + \dots + f\left(\frac{x_{n-1}+x_n}{2}\right) \right)$  or  $w_1 f\left(\frac{x_1+x_2}{2}\right) + w_2 f\left(\frac{x_2+x_3}{2}\right) + \dots + w_{n-1} f\left(\frac{x_{n-1}+x_n}{2}\right)$

Note:  $w = \frac{b-a}{n}$  and applies only for equal sub intervals

14.  $T_n = \frac{w}{2}(y_1 + 2y_2 + \dots + 2y_{n-1} + y_n)$  or  $\frac{1}{2}(w_1(y_1 + y_2) + w_2(y_2 + y_3) + \dots)$

20. Area =  $\int_{x_{left}}^{x_{right}} [f(x)_{top} - f(x)_{down}] dx$  or Area =  $\int_{y_{down}}^{y_{top}} [f(y)_{left} - f(y)_{right}] dy$

21. Vol of rev =  $\pi \int_{x_{left}}^{x_{right}} \left\{ [f(x)_{top} - a]^2 - [f(x)_{down} - a]^2 \right\} dx$  Vol Cross sect. =  $\int_a^b A(x) dx$

**ANTIDIFFERENTIATION (INTEGRATION) RULES**

- $\int x^n dx = \frac{x^{n+1}}{n+1} + C$
- $\int \frac{1}{x} dx = \ln|x| + C$
- $\int \frac{1}{ax+b} dx = \frac{\ln|ax+b|}{a} + C$
- $\int e^{kx} dx = \frac{e^{kx}}{k} + C$
- $\int a^x dx = \frac{a^x}{\ln a} + C$
- $\int \sin kx dx = -\frac{\cos kx}{k} + C$
- $\int \cos kx dx = \frac{\sin kx}{k} + C$
- $\int \sec x \tan x dx = \sec x + C$
- $\int \sec^2 x dx = \tan x + C$
- $\int \csc x \cot x dx = -\csc x + C$

15. **Average Value of f**  $av(f) = f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx$

16. FTC I:  $\int_a^b f'(x) dx = f(b) - f(a)$









17. FTC II i)  $\int_a^x f(t) dt = F(x)$

ii)  $\frac{d}{dx} \int_a^x f(t) dt = f(x)$  iii)  $\frac{d}{dx} \int_{h(x)}^{g(x)} f(t) dt = f(g(x)) \cdot g'(x) - f(h(x)) \cdot h'(x)$

18. Integration by parts:  $\int v du = uv - \int u dv$  use LIPET to select  $u$

19. Integration by substitution:  $\int f(g(x)) g'(x) dx = \int f(u) du$

INTEGRALS (3)

Term	Verbal Description	Symbolic	Graphical
1. Derivative of $f$ at $a$ :	The instantaneous rate of change of the function at $a$ or the slope of the tangent line at $a$	$f'(a) = \left. \frac{df}{dx} \right _{x=a}$ $= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$	
2. Critical Number $c$	A number $c$ in an open $(a, b)$ interval where the derivative is zero or does not exist	$c \in (a, b)$ where $f'(c) = 0$ or $f'(c)$ DNE	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> <math>f'(c) = 0</math>   </div> <div style="text-align: center;"> <math>f'(c)</math> DNE   </div> </div>
3. First Derivative Test	a) If the first derivative changes from <b>negative to positive</b> at $c$ then the function has a <b>relative minimum</b> at $c$ b) If the first derivative changes from <b>positive to negative</b> at $c$ then the function has a <b>relative maximum</b> at $c$	a) If $f'(c)$ $\Delta$ 's from $-$ to $+$ $\Rightarrow f'(c)$ is a min b) If $f'(c)$ $\Delta$ 's from $+$ to $-$ $\Rightarrow f'(c)$ is a max	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">           a)   </div> <div style="text-align: center;">           b)   </div> </div>
4. Concavity Test	a) If the second derivative is <b>positive</b> on an interval $I$ then the function is <b>Concave Up</b> on $I$ b) If the second derivative is <b>negative</b> on an interval $I$ the function is <b>Concave down</b> on $I$	a) If $f''(c) > 0$ on $I$ $\Rightarrow f(x)$ is CU on $I$ b) If $f''(c) < 0$ on $I$ $\Rightarrow f(x)$ is CD on $I$	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>Concave Up <math>f''(x) &gt; 0 \Rightarrow f(x)</math> is CU</p> </div> <div style="text-align: center;">  <p><math>f''(x) &lt; 0 \Rightarrow f(x)</math> is CD Concave Down</p> </div> </div>
5. Point of Inflection at $c$	$f$ : Is a point where the concavity of $f$ changes $f'$ : Is a point where $f'$ changes from increasing to decreasing or decreasing to increasing $f''$ : Is a point where $f''$ changes from positive to negative or negative to positive	$f$ $\Delta$ 's from CU to CD or CD to CU $f'$ $\Delta$ 's from $\nearrow$ to $\searrow$ or $\searrow$ to $\nearrow$ $f''(x)$ $\Delta$ 's from $+$ to $-$ or $-$ to $+$	 <p>Point of Inflection <math>f''(c)</math> <math>\Delta</math>'s from <math>+</math> to <math>-</math> <math>\Rightarrow f(c)</math> is a POI</p>

VOCABULARY (4)

## Motion definitions and Equations

6. Displacement: A Vector quantity that represents the net change in position	$s(t) = x(b) - x(a) = \int_a^b v(t)$	7. Distance: A scalar quantity that represents total movement regardless of sign	$d(t) =  x(b) - x(a)  = \int_a^b  v(t)  dt$
8. Velocity: A Vector quantity that represents the rate of change of position	$v(t) = s'(t)$	9. Speed: A scalar quantity that represents the rate of covering distance	Speed = $ v(t) $
10. Acceleration: A vector quantity that represents the rate of change of velocity	$a(t) = v'(t) = s''(t)$	11. Given initial position $s(a) = C$ the final position is given by $s(b) = s(a) + \int_a^b s'(t) dt$	

## Reciprocal

$$\sin x = \frac{1}{\csc x} \quad \csc x = \frac{1}{\sin x}$$

$$\cos x = \frac{1}{\sec x} \quad \sec x = \frac{1}{\cos x}$$

$$\tan x = \frac{1}{\cot x} \quad \cot x = \frac{1}{\tan x}$$

## Quotient

$$\tan x = \frac{\sin x}{\cos x}$$

$$\cot x = \frac{\cos x}{\sin x}$$


## Pythagorean

$$\sin^2 x + \cos^2 x = 1$$


$$\tan^2 x + 1 = \sec^2 x$$

$$\cot^2 x + 1 = \csc^2 x$$

## Sine Curve



## Cosine Curve



	0	$\pi/6$ (30°)	$\pi/4$ (45°)	$\pi/3$ (60°)	$\pi/2$ (90°)
sin x	0	$1/2$	$\sqrt{2}/2$	$\sqrt{3}/2$	1
cos x	1	$\sqrt{3}/2$	$\sqrt{2}/2$	$1/2$	0
tan x	0	$1/\sqrt{3}$	1	$\sqrt{3}$	Und.
csc x	Und.	2	$2/\sqrt{2}$	$2/\sqrt{3}$	1
sec x	1	$2/\sqrt{3}$	$2/\sqrt{2}$	2	Und.
cot x	Und.	$2/\sqrt{3}$	1	$1/\sqrt{3}$	0

